Ask your Health Care Professional to fill out this form to ensure that you have the proper mixing instructions.

Baby's Name: Today's Date:

Sometimes premature babies require customized feeding strategies. Please follow the directions below for your growing baby.

Caloric Concentration [*]	Water (fl oz)	Scoops [†] of NeoSure Powder (unpacked level scoops)	Approximate Yield (fl oz)
20 Cal/fl oz	4.5	2	5
22 Cal/fl oz (standard dilution)	2	1	2
24 Cal/fl oz	5.5	3	6
27 Cal/fl oz	8	5	9

Health Care Professional Feeding Recommendation:



Health Care Professional Signature

For Home Delivery, call 1-800-FORMULA (1-800-367-6852), Monday-Friday, 8am-5pm CST or visit www.SimilacStore.com for convenient delivery of Similac Expert Care NeoSure product.

* The use of hypo- and hypercaloric formulas should be directed by your baby's doctor. 27 Cal/fl oz or greater formula may not supply enough water for some infants. Hydration status should be monitored and water supplied from other sources if necessary. NOTE: For improved tolerance, it is best to increase by 2 to 4 Cal/fl oz increments.



How to Prepare Similac Expert Care® NeoSure® Powdered Infant Formula

GETTING READY

YOU MUST ADD WATER



Wash your hands, surfaces and utensils.



Ask your baby's doctor if you need to boil (sterilize) utensils, bottles, nipples, rings and water for mixing before use.



If water has been boiled, let it cool to room temperature.

MIXING GUIDE (STANDARD DILUTION)



Pour 4 fl oz of water into clean bottle. Add 2 unpacked level scoops of NeoSure, using only the scoop in can, and return dry scoop to can. Pour 6 fl oz of water into clean bottle. Add 3 unpacked level scoops of NeoSure, using only the scoop in can, and return dry scoop to can.

STORAGE



Cover opened can and store at room temperature. Once mixed, store bottles immediately in refrigerator and feed to baby within 24 hours.

Important:

BE SURE TO USE SCOOP PROVIDED IN CAN.

Store unopened can at room temperature; avoid extreme temperatures.

Use opened can contents within 1 month.

FEEDING BABY



Cap bottle; shake well; attach nipple.



Feed baby.



WARNING: Although powdered infant formulas are not sterile, they may be fed to premature infants or infants who might have immune problems **only as directed by your baby's doctor. Never use a microwave oven to warm formula.** Serious burns can result.



© 2012 Abbott Laboratories 86357/43449/December 2012 LITHO IN USA